

Evaluation of platelet indices in acute, subacute and, chronic phases of deep venous thrombosis

Derin venöz trombozun akut, subakut ve kronik fazlarında platelet indekslerinin değerlendirilmesi

Fatih Ada, Vural Polat

Sivas Cumhuriyet University School of Medicine, Department of Cardiovascular Surgery, Sivas, Turkey


Corresponding author: Fatih Ada, MD, Sivas Cumhuriyet University School of Medicine, Department of Cardiovascular Surgery, Sivas, Turkey


E-mail: drfatihada@gmail.com

Received/Accepted: August 26, 2019 /September 28, 2019

Conflict of interest: There is not a conflict of interest.

SUMMARY

 Fatih Ada

 Vural Polat

Objective: The etiology of deep vein thrombosis disease is multifactorial, and currently, it is seen as a public health problem. Simple, cheap and useful tests that can be used for diagnosis, treatment, and follow-up is increasingly needed day by day because of the increasing number of patients. For this purpose, the changes of platelet indices in routine hemogram tests during DVT follow-up were examined retrospectively.

Method: This study is a retrospective analysis of patients who diagnosed with deep venous thrombosis in Sivas Cumhuriyet University School of Medicine between 01.01.2017-31.12.2018 years. The investigation started after the decision of Sivas Cumhuriyet University Local Ethics Committee No. 2019-02/67. The acute, subacute, and chronic phase parameters of patients with deep venous thrombosis were evaluated. The study included 72 patients who developed DVT. Platelet indexes and other hemogram parameters were analyzed statistically according to the phases of deep vein thrombosis.

Results: 55.6% of the patients were female, and 44.4% were male. DVT was found most commonly in the left femoral region (48.6%). PDW and MPV tended to increase rapidly in the acute phase of deep vein thrombosis and managed to fall in the subacute and chronic phases. However, there was no statistically significant change in platelet count and platelet crit.

Conclusions: PDW and MPV appear to be useful markers in the follow-up of all stages of deep vein thrombosis. It is evident that with the developing technology, more sensitive and specific parameters of platelet indices will be obtained.

Keywords: Deep venous thrombosis, hemogram parameters, platelet indices, mean platelet volume, the platelet distribution width

ORCID IDs of the authors:
F.A. 0000-0002-6953-5906
V.P. 0000-0001-9553-3990

ÖZET

Amaç: Derin ven trombozu etiyolojisi multifaktöriyel bir hastalık olup günümüzde bir halk sağlığı problemi olarak görülmektedir. Artan hasta sayısı ile birlikte tanı, tedavi ve takipte kullanılabilecek basit, ucuz ve kullanışlı testlere ihtiyaç her geçen gün artmaktadır. Bu amaçla rutin hemogram testlerinde yer alan platelet indekslerinin DVT takibinde ki değişimleri retrospektif olarak incelenmiştir.

Yöntem: Bu çalışma, 01.01.2017-31.12.2018 tarihleri arasında Sivas Cumhuriyet Üniversitesi Tıp Fakültesinde, tanı konulmuş derin venöz trombozlu hastaların retrospektif bir analizidir. Çalışmaya, Sivas Cumhuriyet Üniversitesi Yerel Etik Kurulunun 2019-02/67 sayılı kararı sonrası başlandı. Derin venöz trombozlu hastaların akut, subakut ve kronik faz

parametreleri değerlendirildi. Çalışmaya DVT gelişen 72 hasta dâhil edildi. Trombosit indeksleri ve diğer hemogram parametreleri derin ven trombozunun evrelerine göre istatistiksel olarak analiz edildi.

Bulgular: DVT görülen hastaların %55,6'sı kadın ve %44,4'ü erkek idi. En sık sol femoral bölgede (%48,6) DVT saptanmıştır. PDW ve MPV, derin venöz trombozun akut fazında hızlı bir yükselişe geçerken subakut ve kronik fazlarda düşme eğiliminde idi. Bununla birlikte platelet sayısı ve plateletkritin istatistiksel olarak anlamlı bir değişimi yoktu.

Sonuç: Derin ven trombozun tüm evrelerinin takibinde PDW ve MPV iyi birer marker olarak görünmektedir. Gelişen teknoloji ile birlikte platelet indekslerine ait çok daha hassas ve özgül parametrelerin elde edileceği de aşikârdır.

Anahtar sözcükler: Derin venöz tromboz, hemogram parametreleri, trombosit indeksleri, ortalama trombosit hacmi, trombosit dağılım genişliği

INTRODUCTION

Deep venous thrombosis (DVT) has symptoms in a wide range. However, sometimes it can be asymptomatic and defined as subclinical deep venous thrombosis. In addition, It may also lead to pulmonary embolism, and that can cause mortality. Untreated patients may have high morbidity such as; venous ulcer, postphlebotic syndrome. Cancer, postoperative long-term bed dependence, advanced age, obesity, smoking, organ failure, neurological diseases, hereditary causes and increased red blood cell distribution width (RDW) can have a predisposing effect in the etiology of DVT ¹. DVT occurs in Caucasian origin with an incidence of 50-124/100.000 per year ². There is an increasing need for innovations to be used in the prevention, diagnosis, and follow-up for deep venous thrombosis because it is so common high morbidity and mortality disease.

The aim of this study is to examine the changes of platelet indices such as platelet count, platelet crit, mean platelet volume and platelet distribution width in acute, subacute and chronic phases in the follow-up of DVT as thrombosis is the main problem in DVT so that platelet-related studies are frequently encountered ³. In this study, we investigated the relationship between platelet indices of all stages of the DVT process (acute, subacute, chronic).

MATERIAL AND METHODS

In this study patients diagnosed as DVT with doppler ultrasonography between 01.01.2017 and 31.12.2018 at Sivas Cumhuriyet University included and were retrospectively evaluated . The study was started after obtaining permission from the local ethics committee of Sivas Cumhuriyet University numbered 2019-02/67. The written consent form was obtained from all patients. Demographic data such as age, sex, habits, as well as data on doppler ultrasonography and laboratory results were evaluated. Each patient was classified as acute (first 2 weeks), subacute (2-4 weeks), and chronic (after 4 weeks). The course of the platelet indices, as well as the white blood cell count,

hemoglobin level, and the percentage of hematocrit, were analyzed. The patients bigger than 18 years of age as diagnosed of deep vein thrombosis with venous Doppler ultrasonography which complete blood count analysis test was performed in Sivas Cumhuriyet University Hemogram device, were included study. Patients who did not have a hemogram test in all three periods, patients who had a diagnosis and had their hemogram test performed at the external center, and patients under 18 years of age were excluded from the study.

Statistical Analysis:

Statistical analysis was performed using the SPSS (version 24.0, StataCorp LP, College Station, TX, USA) software package. Continuous variables were expressed as mean \pm standard deviation (mean \pm SD), and categorical variables were expressed as a percentage (%). One-Way Anova test was used to test homogeneity of the variables. According to this test result, the Tukey test was performed. However, the Games-Howell test was conducted as the homogeneity of the variables of MPV values was $P < 0.05$. P value $< .05$ was considered as statistically significant in Tukey and Games-Howell tests.

RESULTS

Between 01.01.2017 and 31.12.2018, 481 patients were diagnosed as DVT by doppler ultrasonography at Sivas Cumhuriyet University. However, 72 patients who met inclusion criteria were included in the study. 55,6% of the patients were female, 44,4% were male, and the mean age was $56,2 \pm 20,3$ years. In it's etiology, cases with cancer were seen as 27,8% (n=20), while postpartum DVT was as 9,7% (n=7) (Table 1). DVT seen most commonly in the left femoral region and less in the right iliac region in the lower extremity (Table 2). The DVT rates were equal in the upper extremity. In hemogram parameters, PDW showed a statistically significant difference in all stages of DVT. MPV was significant between acute and chronic stages. No statistically significant difference was found in any of the other parameters (Table 3).

Table 1: Demographic Data of the patients. (DVT: Deep venous thrombosis)

Demographic Data	Percentage (%) or Mean (\bar{x}+sd)	(n) or min-max
Male	44,4	32
Female	55,6	40
Age	56,2±20,3	18-89
Tobacco use	37,5	27
Diabetes Mellitus	34,7	25
Hypertension	40,3	29
Chronic renal disease	16,7	12
Chronic obstructive pulmonary disease	22,2	16
Surgical history	34,7	25
Oral contraceptive use	13,9	10
Cancer	27,8	20
Genetic predisposition	16,7	12
Postpartum DVT	9,7	7

Table 2: Region of the DVT. (DVT: Deep venous thrombosis)

Region of the DVT	Percentage (%)	(n)
Left popliteal	15,3	11
Left femoral	48,6	35
Left iliac	4,2	3
Right popliteal	8,3	6
Right femoral	18	13
Right iliac	2,7	2
Left axillary	1,4	1
Right axillary	1,4	1

Table 3: The result of hemogram parameters. (A-S: Acute-Subacute, A-C: Acute-Chronic, S-C: Subacute-Chronic, SD: standard deviation)

Parameters	Acute Phase (n=72)	Subacute Phase (n=72)	Chronic Phase (n=72)	P
	Mean \pm SD			
White blood cell (WBC$\times 10^9$ /μl)	11,1507 \pm 6.85	10,4967 \pm 15.6	10,5922 \pm 19.1	A-S 0,962>0,05 A-C 0,972>0,05 S-C 0,999>0,05
Hemoglobin (g/dL)	12,6917 \pm 2.49	12,3183 \pm 2.44	12,7014 \pm 2.41	A-S 0,632>0,05 A-C 1,000>0,05 S-C 0,617>0,05
Hematocrit (%)	39,0250 \pm 7.22	38,2111 \pm 6.92	39,2375 \pm 6.99	A-S 0,768>0,05 A-C 0,982>0,05 S-C 0,658>0,05
Platelet (PLT$\times 10^9$ /μl)	221,8347 \pm 107	266,5972 \pm 136	247,9722 \pm 107	A-S 0,062>0,05 A-C 0,382>0,05 S-C 0,612>0,05
Plateletcrit (PCT %)	0,2134 \pm 0.09	0,2448 \pm 0.12	0,2291 \pm 0.10	A-S 0,179>0,05 A-C 0,647>0,05 S-C 0,648>0,05
Mean Platelet Volume (MPV fL)	9,7778 \pm 1.29	9,3083 \pm 1.19	9,2486 \pm 0.96	A-S 0,065>0,05 A-C 0,017<0,05 S-C 0,942>0,05
Platelet Distribution Width (PDW %)	16,3361 \pm 0.43	15,9597 \pm 0,37	15,7153 \pm 0.38	A-S 0,000<0,05 A-C 0,000<0,05 S-C 0,001<0,05

DISCUSSION

Nowadays, deep vein thrombosis is accepted as a public health problem. Venous thromboembolism or its most common form DVT is a significant cause of mortality and morbidity⁴. This disease, which occurs at a rate of 1% in older ages, is most commonly seen in lower extremities and pelvic veins in developed countries⁵. Platelet activation is essential for the development of thrombosis. MPV and PDW increases during platelet activation. However, PDW is more specific than MPV in demonstrating platelet activation⁶. Indeed, the data we obtained from our study supports this. The platelet count begins to decrease due to platelet use during thrombosis process. This promotes the stimulation of bone marrow megakaryocytes by thrombopoietin and accelerates the production of hyperlobulated, more giant platelets with higher DNA content. Therefore, platelets with high MPV are seen in destructive thrombocytopenia because of megakaryocytic stimulation⁷. In thrombocytopenic patients, the splenic pool undergoes destruction and thus an increase in PDW⁸. The rise in PDW and MPV may be independent of platelet count. Indeed, in patients with myeloproliferative thrombocytosis, PDW was

found to be over 17%. Especially in this patient group, PDW value can be observed as high, while the platelet count is normal⁹. Osselaer and colleagues attributed this increase to changes in platelet membrane and adenosine nucleotide content¹⁰.

It is tough to actually make a standardization for the platelet and its indices, which are the subject of many studies. Because there are many parameters that determine the value of this index. For example, the way of blood sampling, temperature, waiting time and the effect of the anticoagulants used so that it is challenging to talk about a standardization¹¹.

There are a lot of studies in the literature on total platelet indices or particularly platelet indices in various disease groups. In addition, indices such as MPV/platelet ratio, neutrophil/lymphocyte ratio, leukocyte/ lymphocyte ratio are used for many different disease groups or treatment methods¹². From these studies, Han and colleagues found that the increased rate of MPV and MPV/Platelet count correlated with DVT². It is a known fact that DVT is associated with many diseases. Wang et al. found that PDW was significantly increased in patients with chronic obstructive pulmonary disease

(COPD) and pulmonary embolism¹³. The source of 70% of pulmonary embolism is DVT¹⁴. However, all DVT doesn't result as pulmonary embolism. Sevük and colleagues reported that the higher MPV and PDW in acute DVT shows a more elevated risk of pulmonary embolism¹⁵.

There are many publications about the use of PDW as a diagnostic or follow-up marker. One of these is the study of Vatankulu and colleagues showed the relationship between PDW elevation and total occlusive coronary artery disease¹⁶. On the other hand, Adam and his colleagues demonstrated the correlation between the PDW elevation and the degree of carotid artery stenosis¹⁷. Li et al. found the high PDW level was associated with increased mortality in melanoma¹⁸. Bhanukumar et al. showed that patients with type 2 DM without a proven history of thrombosis had significantly higher PDW than the control group. They explained the increased amount of thrombosis with PDW elevation and emphasized that PDW value should be considered in diabetic patients for follow-up¹⁹. Sevuk et al. investigated PDW value in different patient groups. They noted that this value decreased in chronic DVT while it had a high value in acute DVT. They emphasized that a reduction in PDW value was a predictor for complete DVT resolution²⁰. Bolayir and colleagues in study of the place of MPV, PDW and platelet / lymphocyte ratio in cerebral venous thrombosis, they observed that MPV and PDW increased significantly in thrombosis group²¹. Kamisli et al. also reported that PDW and MPV values had risen considerably in the early phase of cerebral venous sinus thrombosis compared to the control group²².

Fan and colleagues found high rates of MPV and PDW in acute gangrenous appendicitis²³. Almost all of the publications in the literature mention situations associated with increased MPV and PDW. However, Kaplanoglu et al. found a significant relationship between decreased PDW and spontaneous abortion²⁴.

Ege et al. performed coronary angiography for various reasons and showed that PDW values were higher in patients with coronary artery bypass graft and in patients with saphenous vein graft occlusion than in the patent group²⁵. In the study of Akbulut and colleagues, it was shown that MPV value increased and platelet count decreased in patients with acute myocardial infarction²⁶.

Willians and colleagues examined the full blood parameters in groups of patients with DVT and new diseases. Contrary to classical belief, they found PDW increased while MPV was normal in DVT³.

The relationship between MPV increase and thrombosis status is also discussed in the literature. As a matter of fact, Cil et al. have shown that MPV value is high in patients developing DVT. They also found that this elevation correlated with body-mass index, smoking frequency, hematocrit, and platelet count²⁷. Looking for answers to the question of whether complete blood parameters have a correlation in themselves, Wiwanitkit and his colleagues have shown that only RDW and PDW correlate with each other⁴. The first step of diagnostic processes following clinical suspicion is the examination of colored doppler ultrasonography. D-Dimer and fibrinogen are used to support the diagnosis. In the study of Huang et al., high MPV and PDW level were associated with pulmonary embolism. In particular, the simultaneous elevation of D-Dimer and MPV level was found to have 90,9% specificity in pulmonary embolism²⁸. After definitive diagnosis, the underlying acquired and genetic factors should be investigated simultaneously with the treatment¹⁸. Medical, interventional and surgical methods are included in the procedure. It is possible to prevent mortality and morbidity due to DVT with a timely and correct treatment¹⁴. In the follow-up, colored doppler ultrasonography is frequently used.

CONCLUSION

PDW has been found to be a cheap, simple, and useful marker for monitoring all phases of DVT. It can be used as an adjunct test to diagnose acute DVT and to follow the chronic process. In addition, a new PDW elevation may suggest recurrence in patients with chronic DVT. MPV appears to be useful in differentiating between acute and chronic DVT. It does not seem to be significant for subacute DVT.

REFERENCES

1. Cay N, Unal O, Kartal MG, Ozdemir M, Tola M. Increased level of red blood cell distribution width is associated with deep venous thrombosis. *Blood Coagul. Fibrinolysis* 2013;24(7):727-31.
2. Han JS, Park TS, Cho SY, Joh JH, Ahn HJ. Increased mean platelet volume and mean platelet volume/platelet count ratio in Korean patients with deep vein thrombosis. *Platelets* 2013;24(8):590-3.
3. Willans DJ, Mill SC, Ranney EK. Common thrombotic disorders defined by CBC platelet parameters. *Clin. Appl. Thromb. Hemost.* 1995;1(3):188-201.

4. Wiwanitkit, V. Plateletcrit, mean platelet volume, platelet distribution width: its expected values and correlation with parallel red blood cell parameters. *Clin. Appl. Thromb. Hemost.*2004;10(2):175-8.
5. Wakefield TW, Myers DD, Henke PK. Mechanisms of venous thrombosis and resolution. *Arterioscler. Thromb. Vasc. Biol.*2008;28(3):387-91.
6. Vagdatli E, Gounari E, Lazaridou E, Katsibourlia E, Tsikopoulou F, Labrianou I. Platelet distribution width: a simple, practical and specific marker of activation of coagulation. *Hippokratia* 2010;14(1):28-32.
7. Beyan C. Is mean platelet volume a predictive marker in patients with venous thrombosis?. *Clin. Appl. Thromb. Hemost.* 2012;18(6):670-1.
8. Reddy RS, Khan MI, Phansalkar MD. Platelet Distribution Width (PDW) in Thrombocytopenia. *Indian Medical Gazette* May 2015;169-174.
9. Van der Lelie J, Von dem Borne AK. Platelet volume analysis for differential diagnosis of thrombocytosis. *J. Clin. Pathol.* 1986;39(2):129-33.
10. Osselaer JC, Jamart J, Scheiff JM. (1997). Platelet distribution width for differential diagnosis of thrombocytosis. *Clin. Chem.* 1997;43(6):1072-6.
11. Farias MG, Schunck EG, Dal Bó S, de Castro SM. Definition of reference ranges for the platelet distribution width (PDW): a local need. *Clin. Chem. Lab. Med.* 2010;48(2):255-7.
12. Aldemir M, Pektaş MB, Parlar Aİ, Akıcı Ö, Emren SV, Tecer E, et al. L-carnitine supplementation reduces short-term neutrophil-lymphocyte ratio in patients undergoing coronary artery bypass grafting. *Int J Surg* 2015;100(7):1160-8.
13. Wang M, Zhang J, Ji Q, Yang Q, Zhao F, Li W, et al. Evaluation of platelet distribution width in chronic obstructive pulmonary disease patients with pulmonary embolism. *J Biomark.*2016;10(6):587-96.
14. Sarıgül A, Tanyeli Ö. Derin ven trombozunda güncel tedavi yaklaşımları. *Türk Gogus Kalp Dama* 2007;15(4):316-21.
15. Sevik U, Bahadır MV, Altındag R, Baysal E, Yaylak B, Ay N, et al. Value of serial platelet indices measurements for the prediction of pulmonary embolism in patients with deep venous thrombosis. *Ther Clin Risk Manag*2015;11:1243-9.
16. Vatankulu MA, Sonmez O, Ertas G, Bacaksiz A, Turfan M, Erdogan E, et al. A new parameter predicting chronic total occlusion of coronary arteries: platelet distribution width. *Angiology*2014;65(1):60-4.
17. Adam G, Kocak E, Özkan A, Reşorlu M, Çınar C, Bozkaya H, et al. Evaluation of platelet distribution width and mean platelet volume in patients with carotid artery stenosis. *Angiology* 2015;66(4):375-8.
18. Li N, Diao Z, Huang X, Niu Y, Liu T, Liu ZP, et al. Increased platelet distribution width predicts poor prognosis in melanoma patients. *Scientific reports*, 2017;7(1):2970.
19. Bhanukumar M, Ramaswamy PKH, Peddi NK, Menon WB. Mean Platelet Volume and Platelet Distribution Width as Markers of Vascular Thrombosis in Type 2 Diabetes Mellitus. *J Postgrad Med Edu Res* 2016;50(3):127-131.
20. Sevik U, Altındag R, Bahadır MV, Ay N, Demirtas E, Ayaz F. Value of platelet indices in identifying complete resolution of thrombus in deep venous thrombosis patients. *Indian J Hematol Blood Transfus* 2015;31(1):71-6.
21. Bolayır A, Gökçe ŞF. The role of mean platelet volume, platelet distribution width and platelet/lymphocyte ratio in development of cerebral venous thrombosis. *Cumhuriyet Med J* 2017;39(4):683-91.
22. Kamisli O, Kamisli S, Kablan Y, Gonullu S, Ozcan C. The prognostic value of an increased mean platelet volume and platelet distribution width in the early phase of cerebral venous sinus thrombosis. *Clin. Appl. Thromb. Hemost.* 2013;19(1):29-32.
23. Fan Z, Pan J, Zhang Y, Wang Z, Zhu M, Yang B, et al. Mean platelet volume and

- platelet distribution width as markers in the diagnosis of acute gangrenous appendicitis. *Dis Markers*. 2015;2015:542013.
24. Kaplanoglu M, Yuce T, Bulbul M. (2015). Decreased mean platelet volume is associated with the developing stage of fetoplacental unit in spontaneous abortion. *Int J Clin Exp Med*2015;8(7):11301-6.
25. Ege MR, Guray U, Guray Y, Acikgoz S, Demirkan B. Platelet distribution width and saphenous vein disease in patients after CABG. *Herz* 2013;38(2):197-201.
26. Akbulut N, Yıldız M, Yıldız BŞ, Aktaş S, Yaylacı M. Akut Miyokard İnfarktüsülü Hastalarda Ortalama Trombosit Hacmi ve Trombosit Sayısının Değerlendirilmesi. *MN kardiyoloji* 2006;13(3):180-185.
27. Çil H, Yavuz C, İslamoğlu Y, Tekbaş EÖ, Demirtaş S, Atılgan ZA, et al. Platelet count and mean platelet volume in patients with in-hospital deep venous thrombosis. *Clin. Appl. Thromb. Hemost.* 2012;18(6):650-3.
28. Huang J, Chen Y, Cai Z, Chen P. Diagnostic value of platelet indexes for pulmonary embolism. *Am J Emerg Med*2015;33(6):760-3.